arXiv:math-ph/0406046v1 22 Jun 2004

On generalized fractional kinetic equations

R.K. Saxena
Department of Mathematics and Statistics, Jai Narain Vyas University
Jodhpur 342001, INDIA

A.M. Mathai
Department of Mathematics and Statistics, McGill University,
805 Sherbooke Street West, Montreal, CANADA H3A 2K6

H.J. Haubold
Office for Outer Space Affairs, United Nations,
P.O. Box 500, A-1400 Vienna, AUSTRIA

Abstract

In a recent paper, Saxena et al.[1] developed the solutions of three generalized fractional kinetic equations in terms of the Mittag-Leffler functions. The object of the present paper is to further derive the solution of further generalized fractional kinetic equations. The results are obtained in a compact form in terms of generalized Mittag-Leffler functions. Their relation to fundamental laws of physics is briefly discussed.

1 Introduction

The fundamental laws of physics are written as equations for the time evolution of a quantity $X(t), dX(t)/dt = -AX$, where this could be Maxwell's equations or Schrödinger's equation (if A is limited to linear operators), or it could be Newton's law of motion or Einstein's equations for geodesics (if A may also be a nonlinear operator [2,3,27]). The mathematical solution (for linear operators A) is $X(t) = X(0)exp\{-At\}$.

In thermodynamical or statistical applications one is mostly interested in mean values of the quantity $< X(t) >$. In this case, A is a characteristic time scale $A^{-1} = \tau$ in the evolution equation for $< X(t) >$. It then follows that $< X(t) >$ decays exponentially toward equilibrium $< X(t) >=< (X(0) > exp\{-t/\tau\}$.

In 1988, Tsallis [4] generalized the entropic function of Boltzmann-Gibbs statistical mechanics, $s = -\int dx p(x) ln p(x)$, to nonextensive statistical mechanics with $S_q[p] = \{1 - \int dx [p(x)]^q\}/(q-1)$ that leads to q-exponential distributions $p_q(x) \propto [1-(1-q)x^2/kT]^{1/(1-q)}$. Such a distribution reduces to Gaussian distribution for $q = 1$ and for $q = 2$ to a Cauchy-Lorentz distribution, to name two examples. In an attempt to incorporate Lévy distribution into statistical mechanics, Tsallis has also shown that the above distribution becomes a Lévy distribution for $q > 5/3$. Recently, Tsallis [4] used the mathematical simplicity of reaction-type equations, $dX/dY = Y^q$, to emphasize the natural outcome of the above distribution function $p_q(x)$ which corresponds exactly to the solution of the reaction equation of nonlinear type. The solution has power-law behavior. In the following we show that the fractional generalization of the linear reaction-type equation also leads to power-law behavior. In both cases, solutions can be expressed in terms of generalized Mittag-Leffler functions.

2 Generalized Mittag-Leffler function

A generalization of the Mittag-Leffler function [5,6]

$$E_\alpha(z) := \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + 1)}, \tag{1}$$

and its generalized form

$$E_{\alpha,\beta}(z) := \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + \beta)} \quad (\alpha, \beta \in C, Re(\alpha) > 0) \tag{2}$$

was introduced by Prabhakar [7] in terms of the series representation

$$E^{\gamma}_{\alpha,\beta}(z) := \sum_{n=0}^{\infty} \frac{(\gamma)_n z^n}{\Gamma(\alpha n + \beta)(n!)}, \quad (\alpha, \beta, \gamma \in C, Re(\alpha) > 0), \tag{3}$$

where $(\gamma)_n$ is Pochammer's symbol defined by

$$(\gamma)_n = \begin{cases} 1, n=0 \\ \gamma(\gamma+1)...(\gamma+n-1), n \in N \end{cases} \quad \gamma \neq 0 \tag{4}$$

It is an entire function of order $\rho = [Re(\alpha)]^{-1}$ [7]. This function has been studied by Wiman [8,9], Agarwal [10], Humbert [11] and Humbert and Agarwal [12] and several others. Some special cases of (3) are given below:

$$(i) \quad E_{\alpha}(z) = E^{1}_{\alpha,1}(z), \tag{5}$$

$$(ii) \quad E_{\alpha,\beta}(z) = E^{1}_{\alpha,\beta}(z), \tag{6}$$

$$(iii) \ \Phi(\beta,\gamma;z) = \ {}_1F_1(\beta;\gamma;z) = \Gamma(\gamma)E^{\beta}_{1,\gamma}(z), \tag{7}$$

where $\Phi(\beta,\gamma;z)$ is Kummer's confluent hypergeometric function defined in Erdélyi et al. ([13], p. 248, eq.1]). Mellin-Barnes integral representation for the function defined by (3) follows from the integral

$$E^{\gamma}_{\alpha,\beta}(z) = \frac{1}{2\pi\omega\Gamma(\gamma)} \int_{\Omega} \frac{\Gamma(-s)\Gamma(\gamma+s)(-z)^{s}}{\Gamma(\beta+s\alpha)} ds, \tag{8}$$

where $\omega = (-1)^{1/2}$. The contour Ω is a straight line parallel to the imaginary axis separating the poles of $\Gamma(-s)$ at the points $s = \nu \ \ (\nu = 0, 1, 2, \ldots)$ from those of $\Gamma(\gamma + s)$ at the points $s = -\gamma - \nu \quad (\nu = 0, 1, 2, \ldots)$. The poles of the integrand of (8) are assumed to be simple. (8) can be established by calculating the residues at the poles of $\Gamma(-s)$ at the points , $s = \nu \quad (\nu = 0, 1, 2, \ldots,)$. It follows from (8) that $E^{\gamma}_{\alpha,\beta}(z)$ can be represented in the form

$$E^{\gamma}_{\alpha,\beta}(z) = \frac{1}{\Gamma(\gamma)} H^{1,1}_{1,2}\left[-z\Big|^{(1-\gamma,1)}_{(0,1),(1-\beta,\alpha)}\right], (Re(\alpha) > 0; \alpha, \beta, \gamma \in C), \tag{9}$$

where $H^{1,1}_{1,2}(z)$ is the H-function. A detailed account of the theory and applications of the H-function is available from Mathai and Saxena [14]. This function can also be represented by

$$E^{\gamma}_{\alpha,\beta}(z) = \frac{1}{\Gamma(\gamma)} \ {}_1\Psi_1\left[\begin{matrix}(\gamma,1)\\(\beta,\alpha)\end{matrix}; z\right], \tag{10}$$

where ${}_1\Psi_1(z)$ is a special case of Wright's generalized hypergeometric function ${}_p\Psi_q(z)$ [15,16] ; also see, Erdélyi et al. ([13], Section 4.1), defined by

$${}_p\Psi_q\left[\begin{matrix}(a_1,A_1),\ldots,(a_p,A_p)\\(b_1,B_1),\ldots,(b_q,B_q)\end{matrix}; z\right] = \sum_{n=0}^{\infty} \frac{\prod_{j=1}^{p}\Gamma(a_j + A_j n)}{\prod_{j=1}^{q}\Gamma(b_j + B_j n)} \frac{z^n}{(n)!}, \tag{11}$$

where $1+\sum_{j=1}^{q} B_j - \sum_{j=1}^{p} A_j \geq 0$ (equality only holds for appropriately bounded z). When $\gamma = 1$, (9) and (10) give rise to (12) and (13) given below:

$$E_{\alpha,\beta}(z) = {}_1\Psi_1\left[\begin{matrix}(1,1)\\(\beta,\alpha)\end{matrix}; z\right], \tag{12}$$

$$= H_{1,2}^{1,1}\left[-z\left[\begin{matrix}(0,1)\\(0,1),(1-\beta,\alpha)\end{matrix}\right]\right], \tag{13}$$

where $Re(\alpha) > 0; \alpha, \beta, \gamma \in C$.

If we further take $\beta = 1$ in (12) and (13) we find that

$$E_{\alpha}(z) = {}_1\Psi_1\left[\begin{matrix}(1,1)\\(1,\alpha)\end{matrix}; z\right], \tag{14}$$

$$= H_{1,2}^{1,1}\left[-z\Big|_{(0,1),(0,\alpha)}^{(0,1)}\right], \tag{15}$$

for $Re(\alpha) > 0, z \in C$. The following integral gives the Laplace transform of $E_{\alpha,\beta}^{\gamma}(z)$.

$$\int_0^{\infty} e^{-pt} t^{\beta-1} E_{\alpha,\beta}^{\gamma}(at^{\alpha}) dt = p^{-\beta}(1-ap^{-\alpha})^{-\gamma}, \tag{16}$$

where $Re(p) > |a|^{\frac{1}{Re(\alpha)}}, Re(\beta) > 0, Re(p) > 0$, which can be established by means of the Laplace integral

$$\int_0^{\infty} e^{-pt} t^{\rho-1} dt = \frac{\Gamma(\rho)}{p^{\rho}}, Re(p) > 0, Re(\rho) > 0. \tag{17}$$

For $\gamma = 1$, (16) reduces to an elegant formula

$$\int_0^{\infty} e^{-pt} t^{\beta-1} E_{\alpha,\beta}(at^{\alpha}) dt = p^{-\beta}(1-ap^{-\alpha})^{-1}, \tag{18}$$

where $Re(\beta) > 0, Re(p) > 0, |p| > |a|^{\frac{1}{Re(\alpha)}}$. In an attempt to investigate the functions which when fractionally differentiated (of any order) reappear, Hartley and Lorenzo [17] came across a special function of the form

$$F_q[-a,t] = t^{q-1} \sum_{n=0}^{\infty} \frac{(-a)^n t^{nq}}{\Gamma(nq+q)}, Re(q) > 0, \tag{19}$$

$$= t^{q-1} E_{q,q}(-at^q). \tag{20}$$

This function has been studied earlier by Robotnov [18,19] in connection with hereditary integrals for application to continuum mechanics. The Laplace transform of this function is given by

$$L[F_q(a;t)] = \frac{1}{p^q - a}, Re(q) > 0. \tag{21}$$

A generalization of the F-function is presented by Lorenzo and Hartley [20] by means of the following series representation:

$$\begin{aligned} R_{\nu,\mu}[a,c,t] &= \sum_{n=0}^{\infty} \frac{a^n (t-c)^{(n+1)\nu-\mu-1}}{\Gamma[(n+1)\nu-\mu]}, t > c > 0 & (22) \\ &= (t-c)^{\nu-\mu-1} E_{\nu,\nu-\mu}[a(t-c)^{\nu}], t > c > 0. & (23) \end{aligned}$$

The Laplace transform of the R-function is derived by Lorenzo and Hartley [20] in the form

$$L\left\{R_{\nu,\mu}(a,c,t)\right\} = \frac{e^{-cp} p^{\mu}}{p^{\nu} - a}, \quad c \geq 0, \tag{24}$$

where $Re(\nu - \mu) > 0, Re(p) > 0$. When $c = 0$, (23) reduces to

$$L\left\{R_{\nu,\mu}(a,0,t)\right\} = \frac{p^{\mu}}{p^{\nu} - a}, Re(\nu - \mu) > 0, Re(p) > 0. \tag{25}$$

Finally we recall the definition of Riemann-Liouville operator of fractional integration in the form

$${}_0D_t^{-\nu} f(t) = \frac{1}{\Gamma(\nu)} \int_0^t (t-u)^{\nu-1} f(u) du, \tag{26}$$

with ${}_aD_t^0 f(t) = f(t)$ [21,22,23]. The standard kinetic equation, when integrated, yields

$$N_i(t) - N_0 = c\ {}_0D_t^{-1} N(t), \tag{27}$$

where ${}_0D_t^{-1}$ is the standard Riemann-Liouville integral operator. Here it can be mentioned that in the original paper of Haubold and Mathai [24], the number density of the species $i, N_i = N_i(t)$ is a function of time and $N_i(t=0) = N_0$ is the number density of species i at time $t = 0$. If we drop the index i in (27) and replace c by c^{ν}, then the solution of the generalized equation

$$N(t) - N_0 = -c^{\nu}\ {}_0D_t^{-\nu} N(t), \tag{28}$$

is obtained, Haubold and Mathai [24] as

$$N(t) = N_0 \sum_{k=0}^{\infty} \frac{(-1)^k (ct)^{k\nu}}{\Gamma(k\nu + 1)}. \tag{29}$$

By virtue of (1), (29) can be written in a compact form as

$$N(t) = N_0 E_\nu(-c^\nu t^\nu). \tag{30}$$

In the following, we investigate the solution of two more generalized fractional kinetic equations. The results are obtained in a compact form in terms of generalized Mittag-Leffler functions and are suitable for computation. A detailed account of the operators of fractional integration and their applications is available from a recent survey paper of Srivastava and Saxena [25].

3 Generalized fractional kinetic equations

Theorem 1. If $c > 0, \nu > 0, \mu > 0$, then for the solution of the equation

$$N(t) - N_0 t^{\mu-1} E^{\gamma}_{\nu,\mu}[-c^\nu t^\mu] = -c^\nu_0 D_t^{-\nu} N(t), \tag{31}$$

there holds the formula

$$N(t) = N_0 t^{\mu-1} E^{\gamma+1}_{\nu,\mu}(-c^\nu t^\nu). \tag{32}$$

Proof. By the application of convolution theorem of Laplace transform (Erdélyi et al. [26]) we see that (26) can be written as

$$\begin{aligned} L\left\{{}_0D_t^{-\nu} f(t); p\right\} &= L\left\{\frac{t^{\nu-1}}{\Gamma(\nu)}\right\} L\left\{f(t)\right\}, & (33)\\ &= p^{-\nu} F(p), & (34) \end{aligned}$$

where $F(p) = \int_0^\infty e^{-pu} f(u)du, Re(p) > 0$. Projecting the equation (31) to Laplace transform , we obtain

$$\begin{aligned} N(t) = L[N(t); p] &= N_0 \frac{p^{-\mu}[1 + (p/c)^{-\nu}]^{-\gamma}}{[1 + (p/c)^{-\nu}]} \\ &= N_0 p^{-\mu}[1 + (p/c)^{-\nu}]^{-(\gamma+1)}. \end{aligned} \tag{35}$$

On using the formula (16), we find that

$$L^{-1}[p^{-\mu}\left\{1+(p/c)^{-\nu}\right\}^{-(\gamma+1)}] = t^{\mu-1}E^{\gamma+1}_{\nu,\mu}(-c^{\nu}t^{\nu}). \tag{36}$$

The result (32) now readily follows from (36).

If we set $\gamma = 1$ then (n)! is cancelled. Then in view of the formula

$$\beta E^{2}_{\beta,\gamma}(z) = E_{\beta,\gamma-1}(z) + (1-\gamma+\beta)E_{\beta,\gamma}(z), \tag{37}$$

which follows from the definition of $E^{\gamma}_{\alpha,\beta}(z)$ given by (3) ,we arrive at

Corollary 1.1. If $c > 0, \mu > 0, \nu > 0$, then for the solution of

$$N(t) - N_0 t^{\mu-1}E_{\nu,\mu}[-c^{\nu}t^{\nu}] = -c^{\nu}\ {}_0D_t^{-\nu}N(t), \tag{38}$$

there holds the formula

$$N(t) = \frac{N_0 t^{\mu-1}}{\nu}[E_{\nu,\mu-1}(-c^{\nu}t^{\nu}) + (1-\mu+\nu)E_{\nu,\mu}(-c^{\nu}t^{\nu})]. \tag{39}$$

When $\gamma = 2$, then by virtue of the following identity

$$\begin{aligned} E^{3}_{\beta,\gamma}(z) &= \frac{1}{2\beta^2}[E_{\beta,\gamma-2}(z) + (3\beta+3-2\gamma)E_{\beta,\gamma-1}(z) \\ &+ \left\{2\beta^2+\gamma^2+3\beta-2\gamma-3\beta\gamma+1\right\}E_{\beta,\gamma}(z)], \end{aligned} \tag{40}$$

which follows as a consequence of the definition (3) , we obtain

Corollary 1.2. If $c > 0, \nu > 0, \mu > 0$, then for the solution of

$$N(t) - N_0 t^{\mu-1}E^{2}_{\nu,\mu}[-c^{\nu}t^{\nu}] = -c_0^{\nu}D_t^{-\nu}N(t), \tag{41}$$

there holds the relation

$$N(t) = N_0 t^{\mu-1}E^{3}_{\nu,\mu}(-c^{\nu}t^{\nu}), \tag{42}$$

$$\begin{aligned} &= \frac{N_0 t^{\mu-1}}{2\nu^2}\left[E_{\nu,\mu-2}(-c^{\nu}t^{\nu}) + \left\{3(\nu+1)-2\mu\right\}E_{\nu,\mu-1}(-c^{\nu}t^{\nu})\right. \\ &+ \left.\left\{2(\nu^2+\mu^2)+3\nu-2\mu-3\nu\mu+1\right\}E_{\nu,\mu}(-c^{\nu}t^{\nu})\right]. \end{aligned} \tag{43}$$

Next, if we set $\gamma = 0$, then by virtue of the identity $E^{0}_{\nu,\mu}(z) = \frac{1}{\Gamma(\mu)}$, we arrive at another result given by Saxena et al. [1].

Theorem 2. If $c > 0, b \geq 0, Re(p) > 0, \nu > \mu + 1$, then for the solution of

$$N(t) - N_0 R_{\nu,\mu}(-c^\nu, b, t) = -c^\nu \, {}_0D_t^{-\nu} N(t), \tag{44}$$

there holds the formula

$$N(t) = \frac{N_0}{\nu}(t-b)^{\nu-\mu-1}[E_{\nu,\nu-\mu-1}(-c^\nu(t-b)^\nu) + (\mu+1)E_{\nu,\nu-\mu}(-c^\nu(t-b)^\nu)]. \tag{45}$$

Proof. Taking Laplace transform of both sides of (45), it gives

$$\begin{aligned}
N(t) &= L\{N(t);p\} = N_0 L^{-1}\left[\frac{e^{-bp}p^{\mu-\nu}}{\{1+(c^\nu/p^\nu)\}^2}\right] \\
&= N_0 L^{-1}\left[e^{-bp}p^{\mu-\nu}\sum_{n=0}^{\infty}\frac{(2)_n}{(n)!}(\frac{-c^\nu}{p^\nu})^n\right] \\
&= N_0\sum_{n=0}^{\infty}\frac{(2)_n(-c)^{n\nu}}{(n)!}L^{-1}[e^{-bp}p^{\mu-\nu-n\nu}] \\
&= N_0(t-b)^{\nu-\mu-1}\sum_{n=0}^{\infty}\frac{(2)_n(-c)^{n\nu}(t-b)^{n\nu}}{(n)!\Gamma(n\nu+\nu-\mu)} \\
&= \frac{N_0}{\nu}(t-b)^{\nu-\mu-1}\sum_{n=0}^{\infty}\frac{(-c)^{n\nu}[(n\nu+\nu-\mu-1)+(1+\mu)](t-b)^{n\nu}}{\Gamma(n\nu+\nu-\mu)} \\
&= \frac{N_0}{\nu}(t-b)^{\nu-\mu-1}\left[\sum_{n=0}^{\infty}\frac{\{-c^\nu(t-b)^\nu\}^n}{\Gamma(n\nu+\nu-\mu-1)} + (\mu+1)\sum_{n=0}^{\infty}\frac{\{-c^\nu(t-b)^\nu\}^n}{\Gamma(n\nu+\nu-\mu)}\right] \\
&= \frac{N_0}{\nu}(t-b)^{\nu-\mu-1}[E_{\nu,\nu-\mu-1}\{-c^\nu(t-b)^\nu\} + (\mu+1)E_{\nu,\nu-\mu}\{-c^\nu(t-b)^\nu\}],
\end{aligned}$$

which is same as (45). This completes the proof of theorem 2. If we set $\mu = 0$, theorem 2 reduces to

Corollary 2.1. If $c > 0, b \geq 0, \nu > 1$, then for the solution of

$$N(t) - N_0 R_{\nu,0}(-c^\nu, b, t) = -c^\nu \, {}_0D_t^{-\nu} N(t), \tag{46}$$

there holds the formula

$$N(t) = \frac{N_0}{\nu}(t-b)^{\nu-1}[E_{\nu,\nu-1}(-c^\nu(t-b)^\nu) + E_{\nu,\nu}(-c^\nu(t-b)^\nu)]. \tag{47}$$

For $b = 0$, theorem 2 yields

Corollary 2.2. If $c > 0, \nu > \mu + 1$, then for the solution of

$$N(t) - N_0 R_{\nu,\mu}(-c^\nu, 0, t) = -c^\nu \ {}_0D_t^{-\nu} N(t), \tag{48}$$

there holds the formula

$$N(t) = \frac{N_0}{\nu} t^{\nu-\mu-1} [E_{\nu,\nu-\mu-1}(-c^\nu t^\nu) + (\mu+1) E_{\nu,\nu-\mu}(-c^\nu t^\nu)]. \tag{49}$$

If we further take $\mu = 0$ then the above corollary reduces to the following result:

If $c > 0, \nu > 1$, then the solution of

$$N(t) - N_0 F_\nu[-c^\nu, t] = -c^\nu D_t^{-\nu} N(t), \tag{50}$$

is given by

$$N(t) = \frac{N_0 t^{\nu-1}}{\nu} [E_{\nu,\nu-1}(-c^\nu t^\nu) + E_{\nu,\nu}(-c^\nu t^\nu)]. \tag{51}$$

References

[1] R.K. Saxena, A.M.Mathai, H.J. Haubold, Astrophys. and Space Sci., 282 (2002) 281.

[2] J. Jorgenson, S. Lang, in: Mathematics Unlimited - 2001 and Beyond, Eds. B. Engquist, W. Schmid, Springer-Verlag, Berlin and Heidelberg 2001, 655.

[3] R. Hilfer, in : Applications of Fractional Calculus in Physics, Ed. R. Hilfer, World Scientific, Singapore 2000, 1.

[4] C. Tsallis, in: Nonextensive Entropy: Interdisciplinary Applications, Eds. M. Gell-Mann, C. Tsallis, Oxford University Press, New York 2003,1.

[5] G.M. Mittag-Leffler, C.R. Acad. Sci. Paris (Ser.II)137 (1903) 554.

[6] G.M. Mittag-Leffler, Acta Math. 29 (1905) 101.

[7] T.R. Prabhakar, Yokohama Math. J. 19 (1971) 7.

[8] A. Wiman, Acta Math. 29 (1905) 191.

[9] A. Wiman, Acta Math. 29 (1905) 217.

[10] R.P. Agarwal, C.R. Acad. Sci. Paris 236 (1953) 2031.

[11] P. Humbert, C.R. Acad. Sci. Paris 236 (1953) 1467.

[12] P. Humbert, R.P. Agarwal, Bull. Sci. Math. (Ser.II) 77 (1953) 180.

[13] A. Erdélyi, W. Magnus, F. Oberhettinger, F.G. Tricomi, Higher Transcendental Functions, Vol. 1, McGraw-Hill, New York, Toronto and London 1953.
[14] A.M. Mathai, R.K. Saxena, The H-function with Applications in Statistics and Other Disciplines, Halsted Press, John Wiley & Sons, New York-London-Sydney-Toronto 1978.
[15] E.M. Wright, J. London Math. Soc. 10 (1935) 286.
[16] E.M. Wright, Proc. London Math. Soc. 46 (1940) 389.
[17] T.T. Hartley, C.F. Lorenzo, NASA 1999/ TP-1998-208693 (1998) 1.
[18] Y.N. Robotnov, Tables of a Fractional Exponential Function of Negative Parameters and its Integral (in Russian), Nauka, Russia 1969.
[19] Y.N. Robotnov, Elements of Hereditary Solid Mechanics (in English), MIR Publishers, Moscow 1980.
[20] C.F. Lorenzo, T.T. Hartley, NASA / TP-1999-209424 (1999) 1.
[21] K.B. Oldham, J. Spanier, The Fractional Calculus: Theory and Applications of Differentiation and Integration to Arbitrary Order, Academic Press, New York 1974.
[22] S.G. Samko, A.A. Kilbas, O.I. Marichev, Fractional Integrals and Derivatives. Theory and Applications, Gordon and Breach, Sci. Publ., New York 1990.
[23] K.S. Miller, B. Ross, An Introduction to the Fractional Calculus and Fractional Differential Equations, John Wiley & Sons, New York 1993.
[24] H.J. Haubold, A.M. Mathai, Astrophys. Space Sci. 327 (2000) 53.
[25] H.M. Srivastava, R.K. Saxena, Appl. Math. Comput. 118 (2001) 1.
[26] A. Erdélyi, W. Magnus, F. Oberhettinger, F.G. Tricomi, Tables of Integral Transforms, Vol. 1, McGraw-Hill, New York, Toronto and London 1954.
[27] R. Metzler, J. Klafter, Phys. Rep. 339 (2000) 1.